

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Quantum States Localized on Lagrangian Submanifolds*

François Ziegler (Georgia Southern)

November 8, 2014

*<http://arxiv.org/abs/1310.7882>

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A quantum state

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A quantum state

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state*

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that

- 1 $m(e) = 1$,
- 2 the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbf{C}$ such that

- 1 $m(e) = 1$,
- 2 the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that

- 1 $m(e) = 1$,
- 2 the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that

- 1 $m(e) = 1$,
- 2 the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that

- ① $m(e) = 1$,
- ② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that

- ① $m(e) = 1$,
- ② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

on $\mathbb{C}[G] = \{\text{functions } G \rightarrow \mathbb{C} \text{ with finite support}\}$, is positive.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h)$$

on $\mathbb{C}[G] = \{\text{functions } G \rightarrow \mathbb{C} \text{ with finite support}\}$, is positive.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

Gives rise to unitary G -module $\text{GNS}_m \ni \varphi$ such that $m(g) = (\varphi, g\varphi)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

Gives rise to unitary G -module $\text{GNS}_m \ni \varphi$ such that $m(g) = (\varphi, g\varphi)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

Gives rise to unitary G -module $\text{GNS}_m \ni \varphi$ such that $m(g) = (\varphi, g\varphi)$.
(Put $(\cdot, \cdot)_m$ on $\mathbb{C}[G]$, divide out null vectors and complete; $\varphi = [\delta^e]$.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$ such that

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbb{N}$, $c_j \in \mathbb{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j, \zeta)$.

State of a group G : function $m : G \rightarrow \mathbb{C}$ such that ❶ $m(e) = 1$,
❷ the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\zeta))$.

State of a group G : function $m : G \rightarrow \mathbf{C}$ such that ❶ $m(e) = 1$,
❷ the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\zeta))$.

State of a group G : function $m : G \rightarrow \mathbf{C}$ such that ① $m(e) = 1$,
② the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

State of a group G : function $m : G \rightarrow \mathbf{C}$ such that ❶ $m(e) = 1$,
❷ the sesquilinear form

$$(c, d)_m := \sum_{g, h \in G} \bar{c}_g d_h m(g^{-1}h) \gg 0.$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A *quantum state* is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

- A **quantum representation** (of $\text{Aut}(L)$, for X) is a unitary $\text{Aut}(L)$ -module \mathcal{H} s.t. $m(g) = (\varphi, g\varphi)$ is quantum \forall unit $\varphi \in \mathcal{H}$.

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

- A **quantum representation** (of $\text{Aut}(L)$, for X) is a unitary $\text{Aut}(L)$ -module \mathcal{H} s.t. $m(g) = (\varphi, g\varphi)$ is quantum \forall unit $\varphi \in \mathcal{H}$.
- **Theorem** (Souriau). m quantum $\Rightarrow \text{GNS}_m$ quantum.

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

Examples

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

Examples

None.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

Examples

None. (Unless X is zero-dimensional.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

Examples

None. (Unless X is zero-dimensional.)

Remark. X is a coadjoint orbit of $\text{Aut}(L)$. We might more modestly ask for states and representations of smaller groups (of which X is a coadjoint orbit).

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

(L, ϖ) : Kostant-Souriau line bundle over symplectic manifold (X, ω) .

Definition (Souriau 1990)

A **quantum state** (of $\text{Aut}(L)$, for X) is a state m of $\text{Aut}(L)$ such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{iH_j(x)} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and *complete, commuting* $Z_j \in \text{aut}(L)$ with hamiltonians H_j : $H_j(x) = \varpi(Z_j(\xi))$.

Examples

None. (Unless X is zero-dimensional.)

Remark. X is a coadjoint orbit of $\text{Aut}(L)$. We might more modestly ask for states and representations of smaller groups (of which X is a coadjoint orbit).

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X: coadjoint orbit of a connected Lie group G.

Definition (Souriau 1990)

A **quantum state** (of G, for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbb{N}$, $c_j \in \mathbb{C}$ and commuting $Z_j \in \mathfrak{g}$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

Examples

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A *quantum state* (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

Examples

Too many.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A **quantum state** (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

Examples

Too many. (Unless X is zero-dimensional.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A **quantum state** (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$

for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

Examples

Too many. (Unless X is zero-dimensional.)

- If $X = \{x\}$ is an integral point-orbit, then the unique quantum state for X is the character $m(\exp(Z)) = e^{i\langle x, Z \rangle}$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

X : coadjoint orbit of a connected Lie group G .

Definition (Souriau 1990)

A **quantum state** (of G , for X) is a state m of G such that

$$\left| \sum_{j=1}^n c_j m(\exp(Z_j)) \right| \leq \sup_{x \in X} \left| \sum_{j=1}^n c_j e^{i\langle x, Z_j \rangle} \right|$$


for all choices of $n \in \mathbf{N}$, $c_j \in \mathbf{C}$ and commuting $Z_j \in \mathfrak{g}$.

Examples

Too many. (Unless X is zero-dimensional.)


- If $X = \{x\}$ is an integral point-orbit, then the unique quantum state for X is the character $m(\exp(Z)) = e^{i\langle x, Z \rangle}$.

Let $\hat{g} :=$ (compact) character group of the *discrete* additive group g .




1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} :=$ (compact) character group of the *discrete* additive group g .



1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group


Let $\hat{g} :=$ (compact) character group of the *discrete* additive group g .
We have a dense inclusion $g^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections



1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group


1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} :=$ (compact) character group of the *discrete* additive group g .
We have a dense inclusion $g^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections



1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} :=$ (compact) character group of the *discrete* additive group \mathfrak{g} .
We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections




Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$,
the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} has its spectral measure

This *spectral measure* is the probability measure μ on $\hat{\mathfrak{a}}$ such that
 $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{\mathfrak{a}}} \chi(Z) d\mu(\chi)$. (Bochner.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} :=$ (compact) character group of the *discrete* additive group \mathfrak{g} .
We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections




Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$,
the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} *has its spectral measure*

This *spectral measure* is the probability measure μ on \hat{a} such that
 $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{a}} \chi(Z) d\mu(\chi)$. (Bochner.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} := (\text{compact})$ character group of the *discrete* additive group \mathfrak{g} .
We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections




Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$,
the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} has its spectral measure

This *spectral measure* is the probability measure μ on \hat{a} such that
 $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{a}} \chi(Z) d\mu(\chi)$. (Bochner.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} := (\text{compact})$ character group of the *discrete* additive group \mathfrak{g} . We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections




Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$, the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} has its spectral measure

This **spectral measure** is the probability measure μ on $\hat{\mathfrak{a}}$ such that $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{\mathfrak{a}}} \chi(Z) d\mu(\chi)$. (Bochner.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} := (\text{compact})$ character group of the *discrete* additive group \mathfrak{g} . We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections




Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$, the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} has its spectral measure concentrated on $bX|_{\mathfrak{a}}$, the projection (in $\hat{\mathfrak{a}}$) of the closure bX of X (in \hat{g}).

This **spectral measure** is the probability measure μ on $\hat{\mathfrak{a}}$ such that $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{\mathfrak{a}}} \chi(Z) d\mu(\chi)$. (Bochner.)

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Let $\hat{g} := (\text{compact})$ character group of the *discrete* additive group \mathfrak{g} . We have a dense inclusion $\mathfrak{g}^* \hookrightarrow \hat{g}$, $x \mapsto e^{i\langle x, \cdot \rangle}$, and projections



Theorem

A state m of G is quantum for $X \Leftrightarrow$ for each abelian $\mathfrak{a} \subset \mathfrak{g}$, the state $m \circ \exp|_{\mathfrak{a}}$ of \mathfrak{a} has its spectral measure concentrated on $\text{b}X|_{\mathfrak{a}}$, the projection (in $\hat{\mathfrak{a}}$) of the closure $\text{b}X$ of X (in \hat{g}).

This **spectral measure** is the probability measure μ on $\hat{\mathfrak{a}}$ such that $(m \circ \exp|_{\mathfrak{a}})(Z) = \int_{\hat{\mathfrak{a}}} \chi(Z) d\mu(\chi)$. (Bochner.)

Why “too many” quantum representations?

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Why “too many” quantum representations?

Because this (‘Bohr’) closure operation b is *drastic*:

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Why “too many” quantum representations?

Because this (‘Bohr’) closure operation b is *drastic*:

Theorem (Howe-Z., [dx.doi.org/10.1017/etds.2013.73](https://doi.org/10.1017/etds.2013.73))

(a) *If G is noncompact simple, every nonzero coadjoint orbit is Bohr dense in $\hat{\mathfrak{g}}$, i.e. $bX = \hat{\mathfrak{g}}$.*

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Why “too many” quantum representations?

Because this (‘Bohr’) closure operation b is *drastic*:

Theorem (Howe-Z., [dx.doi.org/10.1017/etds.2013.73](https://doi.org/10.1017/etds.2013.73))

(a) *If G is noncompact simple, every nonzero coadjoint orbit is Bohr dense in $\hat{\mathfrak{g}}$, i.e. $bX = \hat{\mathfrak{g}}$.*

Corollary

(a) *If G is noncompact simple, **every** unitary representation of G is quantum for **every** nonzero coadjoint orbit (!)*

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Why “too many” quantum representations?

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Because this (‘Bohr’) closure operation b is *drastic*:

Theorem (Howe-Z., [dx.doi.org/10.1017/etds.2013.73](https://doi.org/10.1017/etds.2013.73))

- (a) *If G is noncompact simple, every nonzero coadjoint orbit is Bohr dense in $\hat{\mathfrak{g}}$, i.e. $bX = \hat{\mathfrak{g}}$.*
- (b) *If G is connected nilpotent, every coadjoint orbit is Bohr dense in its affine hull.*

Corollary

- (a) *If G is noncompact simple, **every** unitary representation of G is quantum for **every** nonzero coadjoint orbit (!)*

Why “too many” quantum representations?

Because this (‘Bohr’) closure operation b is *drastic*:

Theorem (Howe-Z., [dx.doi.org/10.1017/etds.2013.73](https://doi.org/10.1017/etds.2013.73))

- (a) *If G is noncompact simple, every nonzero coadjoint orbit is Bohr dense in $\hat{\mathfrak{g}}$, i.e. $bX = \hat{\mathfrak{g}}$.*
- (b) *If G is connected nilpotent, every coadjoint orbit is Bohr dense in its affine hull.*

Corollary

- (a) *If G is noncompact simple, **every** unitary representation of G is quantum for **every** nonzero coadjoint orbit (!)*
- (b) *If G is connected nilpotent and X spans \mathfrak{g}^* (reduce to this case by dividing out $\text{ann}(X)$), a unitary representation of G is quantum for $X \Leftrightarrow$ the center acts in it by the character $\exp(Z) \mapsto e^{i\langle X, Z \rangle}$.*

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- Hope that the much-needed selection will arise by restricting attention to states that extend to the whole $\text{Aut}(L)$.
- Require the Bohr closure to be L -invariant. For results along this line see arxiv.org/abs/1011.5056.
- Take this closure seriously. This step is a new interesting step.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

So Souriau's definition is not restrictive enough. 3 ways to proceed:

1. Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
2. *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
3. Take this closure seriously, because it allows interesting states.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- 1 Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- 2 *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- 3 Take this closure seriously, because it allows *interesting states*:

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is *localized at* $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

We also say that the state is *localized on* $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is *localized at* $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

We also say that the state is *localized on* $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

We also say that the state is **localized on** $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_Y$ is a quantum state for Y .

We also say that the state is **localized on** $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_Y$ is a quantum state for Y .

We also say that the state is **localized on** $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

We also say that the state is **localized on** $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_Y$ is a quantum state for Y .

We also say that the state is **localized on** $\pi^{-1}(Y)$, where π is the projection $X \rightarrow \mathfrak{h}^*$. One knows this set is generically a *coisotropic submanifold* — hence at least half-dimensional, and suitable for localizing a system on. We'll mostly consider $Y = \{\text{pt}\}$.

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

One should expect uniqueness of such a state when $\pi^{-1}(Y)$ is *lagrangian* (half-dimensional):

So Souriau's definition is not restrictive enough. 3 ways to proceed:

- ① Hope that the much-needed selection will arise by restricting attention to states that *extend* to the whole $\text{Aut}(L)$.
- ② *Suppress* the Bohr closure implicit in the definition. For results along this line see arxiv.org/abs/1011.5056.
- ③ Take this closure seriously, because it allows *interesting states*:

Definition

Let $H \subset G$ be a closed subgroup and $Y \subset X|_{\mathfrak{h}}$ a coadjoint orbit of H . A quantum state m for X is **localized at** $Y \subset \mathfrak{h}^*$ if the restriction $m|_H$ is a quantum state for Y .

One should expect uniqueness of such a state when $\pi^{-1}(Y)$ is *lagrangian* (half-dimensional): Weinstein (1982) called attaching state vectors to lagrangian submanifolds the FUNDAMENTAL QUANTIZATION PROBLEM.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to x* if, equivalently,

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to* x if, equivalently,

$\langle \exp(t) \rangle$ is a point orbit of H in X .

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to x* if, equivalently,

* $\{x|_H\}$ is a point-orbit of H in \mathfrak{h}^*

* $\langle \chi|_H, \mathfrak{h} \rangle = 0$

* $\chi|_H$ is a character of H .

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to* x if, equivalently,

- $\{x|_h\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [h, h] \rangle = 0$
- $e^{i\langle x, \log h \rangle}$ is a character of H .

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to* x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log|_H}$ is a character of H .

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is *subordinate to* x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log|_H}$ is a character of H .

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^$, namely*

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^$, namely*

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group G : connected, simply connected nilpotent Lie group, X : coadjoint orbit of G , x : chosen point in X .A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^$, namely*

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group G : connected, simply connected nilpotent Lie group, X : coadjoint orbit of G , x : chosen point in X .A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log|_H}$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^$, namely*

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log|_H}$ (discrete induction).

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^*$, namely

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group G : connected, simply connected nilpotent Lie group, X : coadjoint orbit of G , x : chosen point in X .A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^*$, namely

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group G : connected, simply connected nilpotent Lie group, X : coadjoint orbit of G , x : chosen point in X .A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log}|_H$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^*$, namely

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log}|_H$ (discrete induction).

$\mathfrak{a} \subset \mathfrak{h} \Rightarrow x|_{\mathfrak{a}}$ certain;

G : connected, simply connected nilpotent Lie group,

X : coadjoint orbit of G ,

x : chosen point in X .

A connected subgroup $H \subset G$ is **subordinate to** x if, equivalently,

- $\{x|_{\mathfrak{h}}\}$ is a point-orbit of H in \mathfrak{h}^*
- $\langle x, [\mathfrak{h}, \mathfrak{h}] \rangle = 0$
- $e^{ix \circ \log|_H}$ is a character of H .

Theorem

Let $H \subset G$ be maximal subordinate to $x \in X$. Then there is a unique quantum state for X localized at $\{x|_{\mathfrak{h}}\} \subset \mathfrak{h}^*$, namely

$$m(g) = \begin{cases} e^{ix \circ \log(g)} & \text{if } g \in H, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover $\text{GNS}_m = \text{ind}_H^G e^{ix \circ \log|_H}$ (discrete induction).

$\mathfrak{a} \subset \mathfrak{h} \Rightarrow x|_{\mathfrak{a}}$ certain; $\mathfrak{a} \nmid \mathfrak{h} \Rightarrow x|_{\mathfrak{a}}$ equidistributed in $\hat{\mathfrak{a}}$.

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log}|_H$ (usual induction).

This is

- (a) irreducible $\Leftrightarrow H$ is a *polarization at x* (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) equivalent to $I(x, H)$ if $H \neq H'$ are two polarizations at x .

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log}|_H$ (usual induction).
This is

- (a) irreducible $\Leftrightarrow H$ is a *polarization at x* (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) *equivalent* to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

- 1. Quantum states
- 2. Localized states
- 3. Nilpotent groups
- 4. Compact groups
- 5. Euclid's group

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log|_H}$ (usual induction).
This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) *equivalent* to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log|_H}$ (usual induction).
This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log|_H}$ (usual induction). This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

In contrast:

Theorem

Let $H \subset G$ be subordinate to x . Then $i(x, H) := \text{ind}_H^G e^{ix \circ \log|_H}$ is

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log|_H}$ (usual induction). This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

In contrast:

Theorem

Let $H \subset G$ be subordinate to x . Then $i(x, H) := \text{ind}_H^G e^{ix \circ \log|_H}$ is

(a) irreducible $\Leftrightarrow H$ is maximal subordinate to x .

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log}|_H$ (usual induction). This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

In contrast:

Theorem

Let $H \subset G$ be subordinate to x . Then $i(x, H) := \text{ind}_H^G e^{ix \circ \log}|_H$ is

- (a) irreducible $\Leftrightarrow H$ is maximal subordinate to x ;
- (b) inequivalent to $i(x, H')$ if $H \neq H'$ are two polarizations at x .

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log|_H}$ (usual induction).
This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .

In contrast:

Theorem

Let $H \subset G$ be subordinate to x . Then $i(x, H) := \text{ind}_H^G e^{ix \circ \log|_H}$ is

- (a) irreducible $\Leftrightarrow H$ is **maximal subordinate** to x ;
- (b) **inequivalent** to $i(x, H')$ if $H \neq H'$ are two polarizations at x .

Remark

Kirillov (1962) used $I(x, H) := \text{Ind}_H^G e^{ix \circ \log}|_H$ (usual induction). This is

- (a) irreducible $\Leftrightarrow H$ is a **polarization at x** (: subordinate subgroup such that the bound $\dim(G/H) \geq \frac{1}{2} \dim(X)$ is attained);
- (b) **equivalent** to $I(x, H')$ if $H \neq H'$ are two polarizations at x .


In contrast:

Theorem

Let $H \subset G$ be subordinate to x . Then $i(x, H) := \text{ind}_H^G e^{ix \circ \log}|_H$ is

- (a) irreducible $\Leftrightarrow H$ is **maximal subordinate** to x ;
- (b) **inequivalent** to $i(x, H')$ if $H \neq H'$ are two polarizations at x .


Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$



B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-i\alpha} e^{-i(b(r-c) - \frac{1}{2}b^2(t-e))} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but


Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$



B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia}e^{-i\{b(r-c)-\frac{1}{2}b^2(t-e)\}}\psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$




B and C are maximal subordinate but only C is a polarization.

So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia}e^{-i\{b(r-c)-\frac{1}{2}b^2(t-e)\}}\psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$




B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

• $I(x, B)$ is L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$

• $I(x, C)$ is L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$



B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.


All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

① $I(x, B)$ in L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$

② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t \psi = \frac{1}{2}\partial_r^2 \psi$

③ $i(x, C)$ in strong irreducible representations of G

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$



B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but


① $I(x, B)$ in L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$

② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t\psi = \frac{1}{2}\partial_r^2\psi$

③ $i(x, C)$ in almost periodic solutions, $\text{norm}^2 \lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R |\psi|^2 dr$

④ $i(x, B)$ in ℓ^2 functions

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$




B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

- ① $I(x, B)$ in L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$
- ② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t\psi = \frac{1}{2}\partial_r^2\psi$
- ③ $i(x, C)$ in almost periodic solutions, $\text{norm}^2 \lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R |\psi|^2 dr$
- ④ $i(x, B)$ in ℓ^2 functions

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$




B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi\left(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix}\right)$, but

- ① $I(x, B)$ in L^2 functions of $\left(\begin{smallmatrix} r \\ t \end{smallmatrix}\right)$
- ② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t \psi = \frac{1}{2}\partial_r^2 \psi$
- ③ $i(x, C)$ in almost periodic solutions, $\text{norm}^2 \lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R |\psi|^2 dr$
- ④ $i(x, B)$ in ℓ^2 functions

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$




B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)(\begin{smallmatrix} r \\ t \end{smallmatrix}) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix})$, but

- ① $I(x, B)$ in L^2 functions of $(\begin{smallmatrix} r \\ t \end{smallmatrix})$
- ② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t\psi = \frac{1}{2}\partial_r^2\psi$
- ③ $i(x, C)$ in almost periodic solutions, $\text{norm}^2 \lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R |\psi|^2 dr$
- ④ $i(x, B)$ in ℓ^2 functions

Example: Extended Galilei group $G = \left\{ g = \begin{pmatrix} 1 & b & \frac{1}{2}b^2 & a \\ & 1 & b & c \\ & & 1 & e \\ & & & 1 \end{pmatrix} \right\}$



B and C are maximal subordinate but only C is a polarization.
So $i(x, C)$, $I(x, C)$, $i(x, B)$ are irreducible but $I(x, B)$ is not.

All act by $(g\psi)(\begin{smallmatrix} r \\ t \end{smallmatrix}) = e^{-ia} e^{-i\{b(r-c) - \frac{1}{2}b^2(t-e)\}} \psi(\begin{smallmatrix} r-c-b(t-e) \\ t-e \end{smallmatrix})$, but

- ① $I(x, B)$ in L^2 functions of $(\begin{smallmatrix} r \\ t \end{smallmatrix})$
- ② $I(x, C)$ in L^2 solutions of Schrödinger's equation $i\partial_t\psi = \frac{1}{2}\partial_r^2\psi$
- ③ $i(x, C)$ in almost periodic solutions, $\text{norm}^2 \lim_{R \rightarrow \infty} \frac{1}{2R} \int_{-R}^R |\psi|^2 dr$
- ④ $i(x, B)$ in ℓ^2 functions — no Schrödinger equation needed!

1. Quantum states
2. Localized states
3. Nilpotent groups
- 4. Compact groups**
5. Euclid's group

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Theorem

*Every quantum representation of a compact Lie group G is continuous.
The irreducible with highest weight λ is quantum for the coadjoint orbit
with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.*

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Theorem

*Every quantum representation of a compact Lie group G is continuous.
The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.*

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

Theorem

- If μ is dominant integral, then there is a unique quantum state m for $X = G(\mu)$ localized at $\{\mu|_t\} \subset \mathfrak{t}^*$; GNS_m is the irreducible representation with highest weight μ .*
- If μ is dominant and not integral, then there is no such state.*

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

Theorem

- If μ is dominant integral, then there is a unique quantum state m for $X = G(\mu)$ localized at $\{\mu|_{\mathfrak{t}}\} \subset \mathfrak{t}^*$; GNS_m is the irreducible representation with highest weight μ .*
- If μ is dominant and not integral, then there is no such state.*

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

Theorem

- If μ is dominant integral, then there is a unique quantum state m for $X = G(\mu)$ localized at $\{\mu|_{\mathfrak{t}}\} \subset \mathfrak{t}^*$; GNS_m is the irreducible representation with highest weight μ .*
- If μ is dominant and not integral, then there is no such state.*

1. Quantum
states2. Localized
states3. Nilpotent
groups4. Compact
groups5. Euclid's
group

Theorem

Every quantum representation of a compact Lie group G is continuous. The irreducible with highest weight λ is quantum for the coadjoint orbit with dominant element $\mu \Leftrightarrow \lambda \leq \mu$.

So even for compact G , Souriau's definition does not recover the usual 'orbit method' (which posits $\lambda = \mu$). In contrast we have, with $T \subset G$ a maximal torus:

Theorem

- If μ is dominant integral, then there is a unique quantum state m for $X = G(\mu)$ localized at $\{\mu|_{\mathfrak{t}}\} \subset \mathfrak{t}^*$; GNS_m is the irreducible representation with highest weight μ .*
- If μ is dominant and not integral, then there is no such state.*

Euclid's group $G = \left\{ g = \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} : \begin{matrix} A \in \text{SO}(3) \\ c \in \mathbb{R}^3 \end{matrix} \right\}$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Example: TS^2


G acts naturally and symplectically on the manifold $X \simeq TS^2$ of oriented lines (a.k.a. light rays) in \mathbb{R}^3 . 2-form $_{k,s}$:

$$\omega = k d(u, dr) + s \text{Arcs}_u.$$

The moment map

$$\Phi(u, r) = \begin{pmatrix} r \times ku + su \\ ku \end{pmatrix}$$

makes X into a coadjoint orbit of G .



$$\text{Euclid's group } G = \left\{ g = \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} : \begin{matrix} A \in \text{SO}(3) \\ c \in \mathbb{R}^3 \end{matrix} \right\}$$

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Example: TS^2


G acts naturally and symplectically on the manifold $X \simeq \text{TS}^2$ of oriented lines (a.k.a. light rays) in \mathbb{R}^3 . 2-form $_{k,s}$:

$$\omega = k d\langle u, dr \rangle + s \text{Area}_{S^2}.$$

The moment map

$$\Phi(u, r) = \begin{pmatrix} r \times ku + su \\ ku \end{pmatrix}$$

makes X into a coadjoint orbit of G .



$$\text{Euclid's group } G = \left\{ g = \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} : \begin{matrix} A \in \text{SO}(3) \\ c \in \mathbb{R}^3 \end{matrix} \right\}$$

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Example: TS^2


G acts naturally and symplectically on the manifold $X \simeq TS^2$ of oriented lines (a.k.a. light rays) in \mathbb{R}^3 . 2-form $_{k,s}$:

$$\omega = k d\langle u, dr \rangle + s \text{Area}_{S^2}.$$

The moment map

$$\Phi(u, r) = \begin{pmatrix} r \times ku + su \\ ku \end{pmatrix}$$

makes X into a coadjoint orbit of G .



$$\text{Euclid's group } G = \left\{ g = \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} : \begin{matrix} A \in \text{SO}(3) \\ c \in \mathbb{R}^3 \end{matrix} \right\}$$

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Example: TS^2


G acts naturally and symplectically on the manifold $X \simeq TS^2$ of oriented lines (a.k.a. light rays) in \mathbb{R}^3 . 2-form $_{k,s}$:

$$\omega = k d\langle u, dr \rangle + s \text{Area}_{S^2}.$$

The moment map

$$\Phi(u, r) = \begin{pmatrix} r \times ku + su \\ ku \end{pmatrix}$$

makes X into a coadjoint orbit of G .



Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole

$$\psi(\vec{\theta}) = \left(\frac{1}{2\pi} \right)^{3/2} e^{-\frac{1}{2} \vec{\theta}^2}$$

$$\langle \vec{\theta} | \vec{\theta}' \rangle = \delta^3(\vec{\theta} - \vec{\theta}')$$

$$\langle \vec{\theta} | \vec{\theta}' \rangle = \delta^3(\vec{\theta} - \vec{\theta}')$$


1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section




(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i(k e_3, c)} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$


$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$

The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$


$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle ke_3, c \rangle} & \text{if } Ae_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$


$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } Ae_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$

The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.


1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 0$:

We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$


$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$


$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$

The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.


1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$


$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$


$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle k e_3, c \rangle} & \text{if } A e_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } A e_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$


cyclic vector:

$$\psi(r) = e^{-ikz}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle ke_3, c \rangle} & \text{if } Ae_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } Ae_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$

cyclic vector:


$$\psi(r) = e^{-ikz}$$

$$\psi(r) = \frac{\sin \|kr\|}{\|kr\|}$$


The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 0$:


We have localized states on 3 types of lagrangians:



(a): the tangent space
at the north pole



(b): the zero
section



(c): the equator's
normal bundle

$$(a) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\langle ke_3, c \rangle} & \text{if } Ae_3 = e_3, \\ 0 & \text{otherwise.} \end{cases}$$

$$(b) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \frac{\sin \|kc\|}{\|kc\|}$$

$$(c) \quad m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} J_0(\|kc_\perp\|) & \text{if } Ae_3 = \pm e_3, \\ 0 & \text{otherwise} \end{cases}$$

cyclic vector:

$$\psi(r) = e^{-ikz}$$

$$\psi(r) = \frac{\sin \|kr\|}{\|kr\|}$$

$$\psi(r) = J_0(\|kr_\perp\|)$$

The resulting GNS modules can be realized as various spaces of solutions of Helmholtz's equation $\Delta\psi + k^2\psi = 0$, with G-action $(g\psi)(r) = \psi(A^{-1}(r - c))$.

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states

2. Localized states

3. Nilpotent groups

4. Compact groups

5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$F(r) = (B + iE)(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} B = 0, & \operatorname{curl} B = kB, \\ \operatorname{div} E = 0, & \operatorname{curl} E = kE, \end{cases}$$

with G-action $(gF)(r) = AF(A^{-1}(r - c))$. The cyclic vector is $F(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group

Case $s = 1$ (zero section no longer lagrangian):

The unique quantum state localized on the tangent space (a) becomes

$$m \begin{pmatrix} A & c \\ 0 & 1 \end{pmatrix} = \begin{cases} e^{i\alpha} e^{i\langle k e_3, c \rangle} & \text{if } A = e^{j(\alpha e_3)}, \quad (j(\alpha) = \alpha \times \cdot) \\ 0 & \text{otherwise.} \end{cases}$$

$\text{GNS}_m = \{\ell^2 \text{ sections } b \text{ of the tangent bundle } TS^2 \rightarrow S^2\}$, with G-action $(gb)(u) = e^{\langle u, kc \rangle J} A b(A^{-1}u)$ where $J\delta u = j(u)\delta u$. Putting

$$\mathbf{F}(r) = (\mathbf{B} + i\mathbf{E})(r) := \sum_{u \in S^2} e^{-\langle u, kr \rangle J} (b - iJb)(u)$$

one obtains a Hilbert space of almost-periodic solutions of the reduced Maxwell equations

$$\begin{cases} \operatorname{div} \mathbf{B} = 0, & \operatorname{curl} \mathbf{B} = k\mathbf{B}, \\ \operatorname{div} \mathbf{E} = 0, & \operatorname{curl} \mathbf{E} = k\mathbf{E}, \end{cases}$$

with G-action $(g\mathbf{F})(r) = A\mathbf{F}(A^{-1}(r - c))$. The cyclic vector is $\mathbf{F}(r) = e^{-ikz}(e_1 - ie_2)$.

1. Quantum states
2. Localized states
3. Nilpotent groups
4. Compact groups
5. Euclid's group